

RESEARCH

Open Access

Distorted self-perceived weight status and its associated factors among civil servants in Tamale, Ghana: a cross-sectional study

Victor Mogre^{1*}, Prosper P Mwinlenna² and Jeremiah Oladele²

Abstract

Background: Obesity has been described as an epidemic and a major public health concern globally. Distorted self-perceived weight status can negatively impact an individual's decision to lose weight as well as adoption of healthful weight management attitudes. This study described self-perceived weight status among adults working in civil service departments in Tamale, Ghana, and compared it to their classification based on *WHR*. It also examined associations of distorted self-perceived weight status with weight loss attitudes, socio-demographic variables and knowledge levels on the health effects of overweight and obesity.

Methods: This cross-sectional study was undertaken from January 2011 to July 2011 among a sample of 186 civil servants living in Tamale. Out of the sample, 121 were men and 65 were women. Participants' self-perceived weight status, socio-demographic and weight loss attitudes were assessed by means of a 10-item questionnaire. Participants' waist and hip circumferences were measured with appropriate tools and computed into *waist hip ratio (WHR)* and classified based on WHO classifications.

Results: More than 80% of the participants were aged below 40 years. Generally, 56.5% (n = 105) participants had *normal weight* and 31.2% (n = 58) were *centrally obese*. The proportion of participants being *centrally obese* was higher in women compared to men (p < 0.0001). Forty four percent of the studied population had a distorted self-perceived weight status. Less than 10% of participants self-perceived themselves as overweight/obese, in which over 47% were, in fact, *overweight/obese* as measured by *WHR*. Factors associated with distorted self-perceived weight status were being *overweight/obese* (Crude OR = 97.3; (35.8-264.6; p < 0.0001), aged < 40 years (Crude OR = 2.8; 95% CI = 1.3-6.5; p = 0.0102) and having inadequate knowledge on the health effects of overweight/obesity (Crude OR = 3.7; CI = 1.3-11.0; p = 0.0114). Weight loss attitudes and methods used to lose weight were not significantly associated to self-perceived weight status and *WHR* measured weight status.

Conclusions: Self-perceived and *WHR* measured weight status of participants did not conform. Distorted self-perceived weight status was not associated to weight loss attitudes but to being *overweight/obese*, being younger (<40 years) and having inadequate knowledge on the health effects of overweight/obesity. Educating people on accurate weight perception and the health effects of overweight/obesity should be considered in designing public health strategies to curb the rising prevalence of overweight/obesity and other non-communicable diseases in Ghana.

* Correspondence: mogrevictor@yahoo.com¹Department of Human Biology, School of Medicine and Health Sciences, University for Development Studies, Tamale, Ghana

Full list of author information is available at the end of the article

Background

Obesity has been described as an epidemic, an international problem and a major public health concern globally [1,2]. In the UK, the prevalence of obesity was reported to be 23% among men and 24% among women in 2006 [3]. Over the last decade, recent studies indicate that the prevalence of obesity in the US has increased from 12.0% to 19.8% [4,5] and half of its adult population is now overweight or obese [6]. However, it is no longer a disease of industrialized countries but exist in developing countries as well [7]. An obesity prevalence of 10% has been reported in West Africa [8] and specifically 18% in the Republic of Benin [8]. In Ghana, a study conducted by Owiredo et al., [9] in Kumasi, Ghana, reported the prevalence of central obesity to be 27.9% among sedentary workers. Several studies have revealed that the prevalence of obesity in Ghana is increasing especially among women [10-15].

Though several factors have been attributed to the current situation, including urbanization, globalisation, nutritional transition and sedentary lifestyles [7,13,15-18], this study intended to examine the potential contribution of alterations of self-perceived weight status and objective weight. In the present study, a distorted weight perception occurs when one self-perceives his/her weight to be found in a different category (normal weight, overweight, obese) than would have been if determined by an objective method of measurement [19]. Self-perception of one's weight status may differ from both public health and normative cultural standards. In African countries including Ghana, fatness has been associated with wealth, prosperity, affluence and happiness [20,21]. Mvo et al., [21] reported that among the black South African population, obesity has no negative social connotations.

Distorted self-perceived weight status has been extensively studied elsewhere, though has not been adequately investigated in Ghana [19,22-25]. Several studies have demonstrated a strong association between self-perceived weight status and weight control behaviour, independent of objective weight status [26-30].

This study investigated the weight perception of weight among adults working in civil service departments in Tamale, Ghana and compared to their weight classifications based on *WHR*. Furthermore, the study sought to examine the associations between distorted self-perceived weight status and weight loss attitudes (i.e. having tried to lose weight and methods used) and some demographic variables (sex and age) and knowledge on the health effects of overweight and obesity. Exploring the relationship between self-perceived weight status and objectively measured weight status as well as weight control attitudes will be potentially useful in the holistic approach to help tackle the rising prevalence of obesity and other non-communicable diseases in Ghana.

Methods

Participants

This cross-sectional study was conducted between January and July, 2011 among adult civil servants living and working in civil service departments in Tamale, Ghana. Tamale is the capital city of the Northern region of Ghana. It lies between latitude 9°22'N and longitude 0°50'W covering an area of about 922 km². It is about 600 km north of Accra, the capital city of Ghana. The Civil Service forms part of the Public Service of Ghana [31]. There are 13 civil service departments in Tamale with a workforce of about 2500. Three departments were randomly selected from a list of the 13 civil service departments. Participants were selected using a proportionate random sampling technique that included more participants from larger departments, with the help of random number statistical tables. From each selected department, a proportionately determined number of participants were selected from the list of potential participants in that department. Selected personnel who declined to participate or were unavailable to participate were immediately replaced with others from the department.

The inclusion of participants was voluntary and informed consent was obtained from each participant. Participants were also assured of confidentiality of any data taken. A total of 186 out of the 265 (70.2% participation rate) civil servants working in the three sampled civil service departments participated in the study.

Ethical approval

The study was approved by the Ethics Committee of the University for Development Studies, School of Medicine and Health Sciences, Ghana.

Anthropometric variables

Anthropometric measurements included waist circumference, measured midway between the inferior angle of the ribs and the suprailiac crest. Hip circumference, measured as the maximal circumference over the buttocks in centimetres. Both measurements were measured to the nearest 1 cm using a non-stretchable fibre-glass measuring tape (Butterfly, China). During both measurements, subjects were made to stand upright, with arms relaxed at the side, feet evenly spread apart and body weight evenly distributed in accordance with the WHO expert consultation report on waist circumference and waist-hip ratio (WHO, 2004). All anthropometric measurements were taken in a private room allocated for that purpose in each of the sampled departments. *Waist to Hip Ratio (WHR)* was calculated by dividing the waist circumference (cm) by the hip circumference (cm). Men with *WHR* < 0.90, 0.90–0.99 and ≥ 1.0 were classified as *normal*, *overweight* or *obese* respectively, while women were classified in the same categories on the basis of *WHR* of < 0.80, 0.80–0.84 and ≥ 0.85 [32].

Questionnaire

A 10-item self-administered questionnaire was developed to collect information on self-perceived weight status of participants and participants' methods of losing weight and awareness of the health effects of obesity. Weight perception was assessed using the question: "Do you think your weight is: about the right weight, overweight or obese". Socio-demographic data such as gender, age and educational status were also asked. The questionnaire underwent pilot testing for clarity and comprehensibility with a group of 10 civil servants. Based on the outcome of the pretesting, a few minor changes were made. The questionnaire also had questions on method of weight control. Self-perceived weight status of participants was then compared with their measured WHR.

Statistical analysis

Self-perceived weight status was compared to the measured WHR by cross-tabulation and Kappa statistics of agreement. Univariate analysis using simple logistic regression was used to assess influence of different variables on distorted weight perception. The results were reported as crude odds ratio (OR) at 95% confidence interval (CI). The results were expressed as proportion and compared using Fischer's exact test or χ^2 for trend analysis as appropriate. A level of $p < 0.05$ was considered as statistically significant. GraphPad Prism version 5.00 (GraphPad software, San Diego California USA, <http://www.graphpad.com/>) and SPSS version 18 for windows were used for all the statistical analysis.

Results

General characteristics of the studied participants are presented in Table 1. The study included 121 men and 65 men. By WHR, out of the 186 participants, 56.5% ($n = 105$) had *normal weight*, 12.4% ($n = 23$) were *overweight* and 31.2% ($n = 58$) were *centrally obese*. The prevalence of *central obesity* was significantly higher in women than in men ($p = 0.0312$). On the other hand, significantly more men than women had *normal weight* status ($p = 0.0032$).

While 90.9% of the studied population self-perceived themselves as normal weight, 9.1% self-perceived their weight as overweight/obese. From the studied population, more women (95.4%) than men (88.4%) self-perceived their weight as normal. Generally, 44.1% of the studied population had a distorted self-perceived weight status. There was no significant relationship between participants' self-perceived weight status, classification of self-perceived weight status and gender. Of the studied participants, while 61.3% ($n = 114$) had never tried to lose weight, 38.7% had ever tried to lose weight. Exactly the same proportion of the studied participants who ever tried to lose weight used diet and exercise. While, the proportion of men using exercise to lose

Table 1 General characteristics of the studied population (n = 186)

Variable	Total (n = 186)	Men (n = 121)	Women (n = 65)	P value
Age (years)				
< 40	150 (80.6%)	94 (77.7%)	56 (86.2%)	0.1790
≥ 40	36 (19.4%)	27 (22.3%)	9 (13.8%)	
Educational level				
High	167 (89.8%)	108 (89.3%)	59 (90.8%)	0.8056
Weight status measured by WHR				
Normal weight	105 (56.5%)	78 (64.5%)	27 (41.5%)	0.0032
Overweight	23 (12.4%)	12 (10.0%)	11 (16.9%)	0.1707
Obese	58 (31.2%)	31 (25.6%)	27 (41.5%)	0.0312
Self-Perceived weight status				
Overweight/obese	17 (9.1%)	14 (11.6%)	3 (4.6%)	0.1808
Normal weight	169 (90.9%)	107 (88.4%)	62 (95.4%)	
Classification of Self-Perceived weight status				
Distorted	82 (44.1%)	50 (41.3%)	32 (49.2%)	0.2622
Non-distorted	104 (55.9%)	71 (58.7%)	33 (50.8%)	
Ever tried to lose weight				
Yes	72 (38.7%)	41 (33.9%)	31 (47.7%)	0.0824
No	114 (61.3%)	80 (66.1%)	34 (52.3%)	
Methods used to lose weight				
Diet	36 (50.0%)	15 (36.6%)	21 (67.7%)	0.0167
Exercise	36 (50.0%)	26 (63.4%)	10 (32.3%)	
Health personnel's advised me to lose weight				
Yes	32 (44.4%)	21 (51.2%)	11 (35.5%)	0.2337
No	40 (55.6%)	20 (48.8%)	20 (64.5%)	
Has awareness on obesity and its health effects				
Yes	168 (90.3%)	112 (92.6%)	56 (86.2%)	0.1949
No	18 (9.7%)	9 (7.4%)	9 (13.8%)	

Data are presented as proportion and analyzed using Fischer's exact test.

weight was higher than women, the proportion of women using diet to lose weight was higher than men ($p = 0.0167$). More men (51.2%) than women (35.5%) were advised to lose weight by health personnel, even though the differences were not statistically significant. Of the 186 studied participants, 90.3% said they had awareness on the health effects of obesity with insignificant differences between men and women.

The effect of sex, age, weight loss attitudes and WHR measured weight status on the risk of having a distorted self-perceived weight status is presented in Table 2. Using univariate analysis, having an age below 40 years ($OR = 2.8$; 95% $CI = 1.3-6.5$; $p = 0.0102$) and knowledge on the health effects of obesity ($OR = 3.7$; $CI = 1.3-11.0$; $p = 0.0114$) were the variables that significantly increased the risk of having a distorted self-perceived weight status. In addition, participants who were *overweight/obese*,

Table 2 Univariate analysis of some factors for distortion of weight status

Variable	n/N*	Rate of distortion	OR (CI 95%)	P value
Sex				
Male	50/121	41.3%	Ref	Ref
Female	32/65	49.2%	1.4 (0.8-2.5)	0.3003
Age (years)				
20-39	73/150	48.7%	2.8 (1.3-6.5)	0.0102
40-49	9/36	25.0%		
Ever tried to lose weight				
Yes	34/72	47.2%	1.2 (0.7-2.2)	0.4936
No	48/114	42.1%	Ref	Ref
Has awareness on obesity and its health effects				
Yes	69/168	41.1%	Ref	Ref
No	13/18	91.0%	3.7 (1.3-11.0)	0.0114
WHR Status				
Normal Weight	9/105	8.60%	Ref	Ref
Overweight/Obese	73/81	90.1%	97.3 (35.8-264.6)	< 0.0001

*Number of subjects with Distortion/number of subjects in each category. Ref refers to reference point.

were several folds more likely to have a distorted self-perceived weight status compared to participants measured as normal weight (OR = 97.3; 35.8-264.6; $p < 0.0001$) using the univariate analysis.

A comparison between actual weight status as measured by WHR and self-perceived weight status, weight loss attitudes, awareness and age are presented in Table 3. From the over 90% of participants self-perceiving their weight as normal, 56.8% were, in fact, *normal weight*. And of the less than 10% of participants self-perceiving themselves as overweight/obese, over 47% were, in fact, *overweight/obese*. A poor strength of agreement was observed between self-perceived weight status and WHR (Kappa = 0.014 SE = 0.047, CI = -0.078 to 0.107). Among participants who had a distorted self-perceived weight status, while more than 80% of them were *overweight/obese*, only 9% had *normal weight* ($p < 0.0001$). Weight loss attitudes and methods used to lose weight were not significantly associated to weight status measured by WHR. Concerning the decision to lose weight, while, 34.4% of participants who tried to lose weight were *overweight/obese*, 65.6% had *normal weight* ($p = 0.0992$). By proportion, more participants aged 20-39 compared to participants aged 40-49 were *overweight/obese* ($p = 0.1933$).

Discussion

This cross-sectional study among Ghanaian civil servants showed a *central obesity* prevalence of 31.2%. This is consistent with the findings of Owiredun et al., [9] among Ghanaian sedentary workers and Reddy et al.,

Table 3 Comparison between participants' actual weight status (WHR) to self-perceived weight status, weight loss attitudes and age

Variable	Actual weight as measured by WHR			P value
	Normal (%)	Overweight/obesity (%)	Total (n = 186**)	
Self-perceived Weight status				
Normal weight	96 (56.8%)	73 (43.2%)	169 (90.9%)	0.8014
Overweight/obesity	9 (52.9%)	8 (47.1%)	17 (9.1%)	
Classification of self-perceived weight status				
Distorted	9 (10.9%)	73 (89.0%)	82 (44.1%)	< 0.0001
Non-Distorted	96 (92.3%)	8 (7.7%)	104 (55.9%)	
Ever tried to lose weight				
Yes	39 (54.2%)	33 (45.8%)	72 (38.7%)	0.6507
No	66 (57.9%)	48 (42.1%)	114 (61.3%)	
Methods used to lose weight				
Diet	19 (52.8%)	17 (47.2%)	36 (50.0%)	1.0000
Exercise	20 (55.6%)	16 (44.4%)	36 (50.0%)	
Health personnel's advice to lose weight				
Yes	21 (65.6%)	11 (34.4%)	32 (44.4%)	0.0992
No	18 (45.0%)	22 (55.0%)	40 (55.6%)	
Has awareness on obesity and its health effects				
Yes	95 (56.5%)	73 (43.5%)	168 (90.3%)	1.0000
No	10 (55.6%)	8 (44.4%)	18 (9.7%)	
Age (years)				
< 40	81 (54.0%)	69 (46.0%)	150 (80.6%)	0.1933
≥ 40	24 (66.7%)	12 (33.3%)	36 (19.4%)	

Data are presented as proportion and analyzed using Fischer's exact test. *Row percentage, **Column percentage.

[33] among an Industrial population in India. The prevalence of *central obesity* was significantly higher in women compared to men. To a large extent these findings concur with the studies of Puepet et al., [34] among a non-diabetic population in Nigeria; among an apparently healthy population in Uganda [35] and in several studies conducted in Africa [7,36].

A high prevalence of distorted self-perceived weight status has been found in this study. Previously, higher proportions have been reported in Sub-Saharan Africa and elsewhere. A population based survey from the Seychelles reported that 72% of the studied population did not consider themselves to be overweight/obese [37]. In Morocco, 48.5% ($n = 80$) of men vs. 75.2% ($n = 204$) of women had a false perception of their body-weight status in a cross-sectional study of 436 individuals aged 20 years and above [38]. A cross-sectional survey of a sample of 493 adults in Pakistan reported that more than 70% of *obese* participants did not consider themselves as obese [39].

The high prevalence of distorted self-perceived weight status in the present study could be influenced by the predominant culture, physique of peers, media and one's perception of body weight [40]. It has been shown that when overweight/obesity is a common feature of a cultural group, a strong negative social pressure limits involvement in weight control programmes [41]. In this part of the world, many people generally perceive overweight/obesity to be a sign of affluence. A study among University students in Nigeria showed that most subjects believed that being obese gives respect and a sign of living good [42]. As such participants misconceive themselves to be normal when, in fact they were *overweight/obese*.

In this study, a significant proportion of *overweight/obese* (in both genders) participants did not perceive themselves as overweight/obese. In fact, using univariate analysis, *overweight/obese* participants were several folds at risk of having a distorted self-perceived weight status than their normal weight counterparts. Consistent findings have been reported in a population-based survey of Seychelles adults [37], among adolescents in Nigeria [43] and among overweight/obese adults in the US [44]. Another study by Faber [45] among black women in a South African Rural village showed that 2% of Black women perceived themselves to be overweight, while in fact, 95% were overweight. Consistent with other studies, this study has demonstrated a widespread underestimation of body weight and size, especially among overweight/obese individuals [19,25,46].

In this study, age was found to significantly affect distorted self-perceived weight status. We found that participants aged <40 years (in both genders) had a 2.8 risk of having a distorted self-perceived weight status. Most of them underestimated their weight status to be normal when in fact they were *overweight/obese*. In contrast to our findings Bhanji et al., [39] in a study among a sample of adults in Pakistan found that participants aged >40 years were more likely to misperceive their weight status. Furthermore, a study among male firefighters in America found that underestimating one's weight category furthermore increased with age [47]. The inconsistencies could be attributed to social and cultural differences. In Sub-Saharan Africa being overweight/obese is a sign of good health and well-being and is generally accepted as normal weight. As a result, the younger participants can be said to be more concerned about their weight status and would like to misperceive themselves as normal according to the social and cultural norms.

Another important observation of this study was that having inadequate knowledge on the health effects of overweight and obesity was significantly associated to distorted self-perceived weight status. Participants who had inadequate knowledge had crude odds ratio of 2.7

risk of having distorted weight perception. This suggests that they did not consider overweight and obesity as risk factors for chronic diseases such as diabetes mellitus, hypertension, and coronary heart diseases. This is confirmed in a study by Gorynski et al., [48] in which people with diabetes had a poor weight perception. This is worrying in the sense that the distorted weight perception caused by having inadequate knowledge on the health effects of overweight/obesity could lead to a progressive development of obesity resulting in chronic diseases. Distorted self-perceived weight status is a potentially modifiable risk factor [44] and as such educating people on the health effects of overweight and obesity should be considered in public health interventions in the future.

Even though a high prevalence of *central obesity* has been found in this study, a large proportion of *normal weight* and *overweight/obese* participants had never attempted to lose weight. Participants appeared to be unconcerned about their weight status. In a study of black women, in South Africa few obese participants wanted to or had ever tried to lose weight [23]. Participants' apparent poor attitude towards losing weight could be due to the fact that a large proportion of the participants in this study misperceived themselves to be normal weight, when in fact they were either overweight or obese. Self-defined body weight is very critical, as it plays a role in the weight management practices of individuals as well as their nutrition habits [49]. A study by Duncan et al., [44] reported that overweight/obese men and women who misperceived their overweight status were 60% (RR 0.40, 95% CI 0.30-0.52) and 56% (RR 0.44, 95% CI 0.32-0.59) less likely than those who accurately perceived their overweight status to have attempted weight loss during the past year. The findings of this study and several others suggest that weight perception should be considered in designing weight loss interventions [44,50,51].

When participants' choice of method to lose weight was stratified by gender, more women preferred dieting as a method to lose weight while more men preferred exercise as a method to lose weight. This is consistent with the findings of Rasheed [52] in a study among female students in Saudi Arabia, in which women preferred dieting to exercise to lose weight. According to that study women had a poor attitude towards exercise. In a study in Ghana, which assessed the healthy life behaviours among a sample population, men were found to have a higher physical activity level than women [53]. Several other studies have also reported women having low physical activity levels compared with men, indicating the low preference of women for exercise [7,42,53].

The inability of participants in this study to self-perceive themselves as overweight/obese can hinder the effectiveness of weight control programmes because they

will have less incentives to control weight [45]. Social determinants of weight perception are of clinical and public health significance [45,54] and further research is recommended to explore the macro and micro social processes involved. This will help curb the rising prevalence of overweight/obesity in Ghana.

It is critical to note that this study is subject to some limitations. This study did not consider socio-economic status of the participants which have been shown to influence weight perception. Some questions on socio-economic status were asked but the responses were either missing or ambiguous. Secondly, our study did not consider body size and body image perception which are important factors of weight perception and weight loss attitudes. Thirdly, this study was carried out on a small sample in an urban area and as a result the findings cannot be generalized to the entire Ghanaian population. However, it can serve as a reference point for further studies in the country. Finally, the cross-sectional nature of this study cannot investigate the direction of causality.

Conclusion

A high prevalence of *central obesity* and distorted self-perceived weight status was observed. Weight loss attitudes were not associated to distorted self-perceived weight status. Distorted self-perceived weight status was common among *overweight/obese* participants, younger participants and participants who lacked knowledge on the health effects of overweight and obesity. Weight perception should be considered in designing weight loss intervention in the future. In addition, people should be educated on the health effects of overweight and obesity. These are important strategies that should be considered in addressing the rising prevalence of overweight and obesity in Ghana.

Competing interests

The right of we the authors to examine, analyze, and publish the data of the research is not infringed upon by any contractual agreement or conflict of interest.

Authors' contributions

VM, analysis and Interpretation of data, drafting of the manuscript and critical revision of the manuscript. PPM, concept and design. JO, data acquisition. All authors read and approved the final manuscript.

Acknowledgements

Authors wish to thank Dr. Nafiu Amidu, Editor-in-Chief of the Journal of Medical and Biomedical Sciences and Dr. Alhassan Abass, Lecturer, Department of Human Biology, School of Medicine and Health Sciences, UDS for their professional guidance and encouragement during the write up of this paper.

Author details

¹Department of Human Biology, School of Medicine and Health Sciences, University for Development Studies, Tamale, Ghana. ²Department of Community Nutrition, School of Medicine and Health Sciences, University for Development Studies, Tamale, Ghana.

Received: 26 April 2013 Accepted: 26 October 2013
Published: 6 November 2013

References

- Hussin DK, Mohammad IH, Al-Hamad HA, Makkoul G, Elshazly M: Weight status and perceived body size image in overweight and obese children 8-12 years old. *Alexandria J Med* 2011, **47**:365-371.
- Chang JW, Chen YC, Chen CY, Chen JT, Chen SK, Wang YC: Correlation of genetic instability with mismatch repair protein expression and p53 mutations in non-small cell lung cancer. *Clin Cancer Res* 2000, **6**:1639-1646.
- Allender S, Peto V, Scarborough P, Boxer A, Rayner M: Diet, physical activity and obesity statistics. 2012.
- Mokdad AH, Bowman BA, Ford ES, Vinicor F, Marks JS, Koplan JP: The continuing epidemics of obesity and diabetes in the United States. *JAMA* 2001, **286**:1195-1200.
- Mokdad AH, Serdula MK, Dietz WH, Bowman BA, Marks JS, Koplan JP: The spread of the obesity epidemic in the United States, 1991-1998. *JAMA* 1999, **282**:1519-1522.
- Flegal KM, Carroll MD, Ogden CL, Johnson CL: Prevalence and trends in obesity among US adults, 1999-2000. *JAMA* 2002, **288**:1723-1727.
- Wahab KW, Sani MU, Yusuf BO, Gbadamosi M, Gbadamosi A, Yandutse MI: Prevalence and determinants of obesity-a cross-sectional study of an adult Northern Nigerian population. *Int Arch Med* 2011, **4**:10.
- Abubakari AR, Lauder W, Agyemang C, Jones M, Kirk A, Bhopal RS: Prevalence and time trends in obesity among adult West African populations: a meta-analysis. *Obes Rev* 2008, **9**:297-311.
- Owiredo WKBA, Amidu N, Gockah-Adapoe E, Ephraim RKD: The prevalence of metabolic syndrome among active sportsmen/sportswomen and sedentary workers in the Kumasi metropolis. *J Sci Technol* 2011, **31**:23-36.
- Steyn K, Damasceno A: Lifestyle and related risk factors for chronic diseases. *Disease and mortality in sub-Saharan Africa*. 2006, **2**.
- Kruger HS, Puoane T, Senekal M, van der Merwe MT: Obesity in South Africa: challenges for government and health professionals. *Public Health Nutr* 2005, **8**:491-500.
- Chitson P: Health care implications of non-communicable diseases. In *Proceedings of the IGU conference on health problems*. IGU; 1994:1-9.
- Mogre V, Mwinlenaa PP, Oladele J, Amalaba A: Impact of physical activity levels and diet on central obesity among civil servants in Tamale metropolis. *J Med Biomed Sci* 2012, **1**.
- Aryeetey R, Ansong J: Overweight and hypertension among college of health sciences employees in Ghana. *Afr J Food, Agric Nutr Dev* 2011, **11**:5444-5456.
- Addo J, Smeeth L, Leon DA: Obesity in urban civil servants in Ghana: association with pre-adult wealth and adult socio-economic status. *Public Health* 2009, **123**:365-370.
- Candib LM: Obesity and diabetes in vulnerable populations: reflection on proximal and distal causes. *Ann Fam Med* 2007, **5**:547-556.
- Popkin BM, Gordon-Larsen P: The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord* 2004, **28**(Suppl 3):S2-S9.
- Popkin BM: The nutrition transition and its implications in lower-income countries. *Public Health Nutr* 1997, **1**:5-21.
- Johnson-Taylor WL, Fisher RA, Hubbard VS, Starke-Reed P, Eggers PS: The change in weight perception of weight status among the overweight: comparison of NHANES III (1988-1994) and 1999-2004 NHANES. *Int J Behav Nutr Phys Act* 2008, **5**:9.
- Chesler J: A study of attitudes and knowledge concerning obesity in an urban African community. *S Afr Med J* 1961, **35**:129-131.
- Mvo Z, Dick J, Steyn K: Perceptions of overweight African women about acceptable body size of women and children. *Curationis* 1999, **22**:27-31.
- Kurdak H, Bozdemir N, Saatci E, Ozturk P, Ozcan S, Akpinar E: Self-perceived body weight status and weight-control behaviors of high school students in a southern city of Turkey. *Coll Antropol* 2010, **34**:1295-1302.
- Kruger HS, Van Aardt AM, Walker ARP, Bossman MJ: Obesity in African hypertensive women: problems in treatment. *S Afr J Food Sci Nutr* 1994, **8**:106-112.
- Emslie C, Hunt K, Macintyre S: Perceptions of body image among working men and women. *J Epidemiol Community Health* 2001, **55**:406-407.
- Inoue M, Toyokawa S, Miyoshi Y, Miyano Y, Suzuki T, Suyama Y, Inoue K, Kobayashi Y: Degree of agreement between weight perception and body mass index of Japanese workers: my health up study. *J Occup Health* 2007, **49**:376-381.
- Crawford D, Campbell K: Lay definitions of ideal weight and overweight. *Int J Obes Relat Metab Disord* 1999, **23**:738-745.

27. Forman MR, Trowbridge FL, Gentry EM, Marks JS, Hogelin GC: **Overweight adults in the United States: the behavioral risk factor surveys.** *Am J Clin Nutr* 1986, **44**:410–416.
28. Riley NM, Bild DE, Cooper L, Schreiner P, Smith DE, Sorlie P, Thompson JK: **Relation of self-image to body size and weight loss attempts in black women: the CARDIA study: coronary artery risk development in young adults.** *Am J Epidemiol* 1998, **148**:1062–1068.
29. Strauss RS: **Self-reported weight status and dieting in a cross-sectional sample of young adolescents: national health and nutrition examination survey III.** *Arch Pediatr Adolesc Med* 1999, **153**:741–747.
30. Serdula MK, Collins ME, Williamson DF, Anda RF, Pamuk E, Byers TE: **Weight control practices of US adolescents and adults.** *Ann Intern Med* 1993, **119**:667–671.
31. Addo J, Smeeth L, Leon DA: **Smoking patterns in Ghanaian civil servants: changes over three decades.** *Int J Environ Res Public Health* 2009, **6**:200–208.
32. WHO: **Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies.** *Lancet* 2004, **363**:157–163.
33. Reddy KS, Prabhakaran D, Chaturvedi V, Jeemon P, Thankappan KR, Ramakrishnan L, Mohan BV, Pandav CS, Ahmed FU, Joshi PP, et al: **Methods for establishing a surveillance system for cardiovascular diseases in Indian industrial populations.** *Bull World Health Organ* 2006, **84**:461–469.
34. Puepet FH, Ohwovoriole AE: **Prevalence of risk factors for diabetes mellitus in a non-diabetic population in Jos, Nigeria.** *Niger J Med* 2008, **17**:71–74.
35. Maher D, Waswa L, Baisley K, Karabarinde A, Unwin N, Grosskurth H: **Distribution of hyperglycaemia and related cardiovascular disease risk factors in low-income countries: a cross-sectional population-based survey in rural Uganda.** *Int J Epidemiol* 2011, **40**:160–171.
36. Shayo GA, Mugusi FM: **Prevalence of obesity and associated risk factors among adults in Kinondoni municipal district, Dar es Salaam Tanzania.** *BMC Public Health* 2011, **11**:365.
37. Alwan H, Viswanathan B, Williams J, Paccaud F, Bovet P: **Association between weight perception and socioeconomic status among adults in the Seychelles.** *BMC Public Health* 2010, **10**:467.
38. Lahmam A, Baali A, Hilali MK, Cherkaoui M, Chapuis-Lucciani N, Boetsch G: **Obesity, overweight and body-weight perception in a high atlas Moroccan population.** *Obes Rev* 2008, **9**:93–99.
39. Bhanji S, Khuwaja AK, Siddiqui F, Azam I, Kazmi K: **Underestimation of weight and its associated factors among overweight and obese adults in Pakistan: a cross sectional study.** *BMC Public Health* 2011, **11**:363.
40. Tang J, Yu Y, Du Y, Ma Y, Zhu H, Liu Z: **Association between actual weight status, perceived weight and depressive, anxious symptoms in Chinese adolescents: a cross-sectional study.** *BMC Public Health* 2010, **10**:594.
41. Kumanyika S, Wilson JF, Guilford-Davenport M: **Weight-related attitudes and behaviors of black women.** *J Am Diet Assoc* 1993, **93**:416–422.
42. Ojofeitimi EO, Adeyeye AO, Fadiora AO, Kuteyi AO, Faborode TG, Adegbenro CA, Bakare OE, Setiloane K, Towobola KS: **Awareness of obesity and its health hazard among women in a university community.** *Pak J Nutr* 2007, **6**:502–505.
43. Sabageh OA, Ogunfowokan AA, Ojofeitimi OE: **Obesity and body image discrepancy among school adolescents in Ile-Ife, Osun State, Nigeria.** *Pak J Nutr* 2013, **12**:377–381.
44. Duncan DT, Wolin KY, Scharoun-Lee M, Ding EL, Warner ET, Bennett GG: **Does perception equal reality? Weight misperception in relation to weight-related attitudes and behaviors among overweight and obese US adults.** *Int J Behav Nutr Phys Act* 2011, **8**:20.
45. Faber M, Kruger HS: **Dietary intake, perceptions regarding body weight, and attitudes toward weight control of normal weight, overweight, and obese black females in a rural village in South Africa.** *Ethn Dis* 2005, **15**:238–245.
46. Brener ND, Eaton DK, Lowry R, McManus T: **The association between weight perception and BMI among high school students.** *Obes Res* 2004, **12**:1866–1874.
47. Baur DM, Christophi CA, Tsismenakis AJ, Jahnke SA, Kales SN: **Weight-perception in male career firefighters and its association with cardiovascular risk factors.** *BMC Public Health* 2012, **12**:480.
48. Gorynski P, Krzyzanowski M: **A study of the self-perception of being overweight in adult inhabitants of Cracow.** *J Clin Epidemiol* 1989, **42**:1149–1154.
49. Argani L, Toselli S, Gualdi-Russo E: **Body image and growth in Italy.** *Coll Antropol* 2008, **32**:413–418.
50. Jones M, Grilo CM, Masheb RM, White MA: **Psychological and behavioral correlates of excess weight: misperception of obese status among persons with Class II obesity.** *Int J Eat Disord* 2010, **43**:628–632.
51. Skinner AC, Weinberger M, Mulvaney S, Schlundt D, Rothman RL: **Accuracy of perceptions of overweight and relation to self-care behaviors among adolescents with type 2 diabetes and their parents.** *Diabetes Care* 2008, **31**:227–229.
52. Rasheed P: **Overweight status: body image and weight control beliefs and practices among female college students.** *Ann Saudi Med* 1999, **19**:365–369.
53. Tagoe HA, Dake FA: **Healthy lifestyle behaviour among Ghanaian adults in the phase of a health policy change.** *Global Health* 2011, **7**:7.
54. Chang VW, Christakis NA: **Self-perception of weight appropriateness in the United States.** *Am J Prev Med* 2003, **24**:332–339.

doi:10.1186/2049-3258-71-30

Cite this article as: Mogre et al.: Distorted self-perceived weight status and its associated factors among civil servants in Tamale, Ghana: a cross-sectional study. *Archives of Public Health* 2013 **71**:30.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

